

Earth tides and ocean tides

Because so obvious + important to seafaring nations, ocean tides observed far back into history.

Before Newton many colorful "explanations".

Ch. IV of The Tides (1898) by Sir G. Darwin, charming Victorian book for lay audience:

p. 79, from Koran: displacement by foot of an angel

p. 76, "Chinese writers have suggested... that H_2O is the blood of the $\oplus$ and that the tides are the beating of its pulse".


Many early observers ~~noticed~~ noticed correlation with α , esp. spring and neap tides with full and new and half α , respectively.

Early explanations of this: heat from $\odot$ dries up oceans, moisture from α etc.

Greeks and Romans had little cause to ponder the tides as they are very small everywhere in the Mediterranean.

1687 Newton's Principia : first correct discussion in terms of gravitational attraction of $\odot$, $\oplus$ on oceans. This a static theory. Concept of equilibrium tide still plays important role in theory.

Next important development 1774
Laplace's Mechanique Celeste : realized that problem of ocean tides a dynamical one. Can think of tide as a deep water wave




$\lambda \gg h$, thus
phase velocity
 $c = \sqrt{gH}$

$H \sim 5$ km, deep oceans

$$c \sim (10^3 \cdot 5 \cdot 10^5)^{1/2} \sim 2 \cdot 10^4 - 3 \cdot 10^4 \text{ cm/s} \\ \sim 700 - 1000 \text{ km/hr}$$

Across an ocean basin + return in a time of order 12 hr = period of forcing $\Rightarrow$ resonance effects; this invalidates a quasi-static theory.

Analogy: a pendulum (or any other simple harmonic oscillator)




period $T = 2\pi \sqrt{l/g}$, time to complete one cycle.

Say one pushes laterally with a force which varies sinusoidally with a period $T_{\text{force}} \gg T$. Then response will be quasi-static or equilibrium, i.e. all ~~forces~~ forces will be in static balance at all times, acceleration of mass negligible. But if $T_{\text{force}} \sim T$ resonance effects can occur, response more complicated, must solve eqns of motion with acceleration term $\ddot{x}$

Laplace derived eqns of motion for tides on a sphere including acceleration terms and rotation. Only in last few years have computers been adequate to solve L.T.E. numerically for realistic oceanic depths $h(\theta, \phi)$ on $\oplus$ surface.

Lord Kelvin designed first self-recording tide gauge ~ 1860's, previously measured on tide poles (meter sticks read by eye), difficulty with waves, averaged by eye.

Kelvin's gauge damped out the waves so could record mechanically.



Study of tides long hindered by inability to make measurements in deep oceans, only on coasts + scattered islands.



Two techniques now for measuring deep-sea tides:

1. pressure gauges on bottom: tides are deep water waves so $\Delta p = \rho g h$ on bottom ($h =$ height of tide); technology: free-fall retrievable capsules.

2. radar altimetry from satellites, if can measure δ to $\pm$ few cm, can measure variations, this method holds much promise, few concrete results to date.

Kelvin 1876 also first to point out that tidal forces would cause deformations of solid $\oplus$ as well, since $\oplus$ not completely rigid.


Surface of solid $\oplus$ raised by 10-20 cm, tidal strains $\Delta l/l \sim 10^{-7}$ or 10^{-8} measured with strain meters.



$$\Delta l/l \sim 10^{-7} \text{ to } 10^{-8}$$

Two types in common use:

1. Benioff quartz rod, low coeff. of thermal expansion, still must site in mine or tunnel, typically uses capacitive transducers



Alternative: a stretched

Invar wire


measure $C(t)$ by measuring

$\omega = \frac{1}{\sqrt{LC}}$ of a tuned circuit

typically $l \sim 50-100$ m, the longer the better to average out local "geology". If $l = 100$ m

then $\Delta l \sim 0.01$ to 0.001 mm.

2. laser interferometer: stable and precise, but more expensive, not too many $\exists$. Might look like this:




measures strain in fringes, or wavelengths of He-Ne light.

Two problems: changes in refractive index of air, must evacuate pipe, expensive, pumps, etc. esp. for a ~ 1 km instrument.

under development: 3 "color" system measure temp, humidity of air + $\Delta l/l$ simultaneously.

Second problem: He-Ne laser not sufficiently stable, must be stabilized, e.g. with methane absorption servo cell.




mirror of resonator on (essentially)
a loudspeaker coil

There are also tidal gravity fluctuations
 $\Delta g / g \sim 10^{-7}$ to $10^{-8} \sim 0.1$ to 0.01
 mgal (recall $g \sim 980 \text{ gal} \equiv \text{cm/s}^2$)


Measured with long-period seismometers
 Essentially mass on a spring. Three
 types in use:

1. Lacoste:




This is the
 basic geometry.

For use in DA network, electrostatic
capacitive feedback has been added
 as a way to measure deflection of
 mass; general idea:



apply a
 voltage $V(t)$ to keep paddle centered
 between two plates. Actual "signal"
 is servo voltage required to do this.

2. Block-Moore: capacitive feedback system
 first developed for use with
 this instrument, only the "spring"
 is different.



Here the paddle plays the role of the mass too



3. superconducting : the ultimate in stability, sensitivity and \$.

A superconducting sphere about 2 cm in diameter is levitated in a magnetic field. Liquid He temps. guarantee a very low Brownian noise level. Electrostatic feedback is again used to maintain the posn. of the ball when g changes.


There are also tiltmeters that measure tilt $\equiv$

horizontal acceleration, e.g. liquid level tiltmeters (Michelson-Gale)

Earth tides much more nearly equilibrium than ocean tides, fund. period of solid $\oplus$ ${}_0S_2$ mode ~ 54 min. period $\ll 12$ hr. semi-diurnal tide. Time for P wave to go thru center of $\oplus$ and return ~ 40 min.

$\oplus$ tide measurements are however contaminated by ocean loading.


C



causes extra strains, especially near coasts.

This makes it very difficult to determine some things we'd like to know, e.g. fraction of tidal energy dissipated by bodily friction.

Why are there 2 tides / day instead of one?
 Grav. attraction of C at pts. on $\oplus$ surface (i.e. on ocean H_2O particles) varies from place to place.
 Neglect $\odot$ for moment.



↑ one per month
 C

Looks like should only be 1 tide / day as $\oplus$ rotates under tidal bulge (by day we mean lunar day)

But actually $\oplus$ and $\ominus$ revolve mutually about their common c.o.m.

Every pt. on $\oplus$ experiences the same orbital acceleration. This point obscured in many if not most elementary discussions like this one.

Where is c.o.m. ? $\oplus \ominus$ - distance
 $\sim 60 \oplus$ radii, $M_{\ominus} / M_{\oplus} = 1/81.3$.
 Thus:




$$81.3x = 60 - x$$

$$x = \cancel{60} / 82.3$$

$$\approx 3/4$$

$\oplus \ominus$ c.o.m. about $3/4$ of way out to $\oplus$ surface.




motion of every pt. on $\oplus$ is superposition of orbital accel. of c.o.m. $\leftarrow$ is not a centripetal acceleration + $\oplus$ rotation about c.o.m. but an orbital acceleration of same length (as well see in derivation below)

it's the same at every pt. on $\oplus$ because every pt. experiences the same orbital motion, hence same $\frac{d^2}{dt^2}$ motion

Resultant looks like: tries to squeeze $\oplus$ into an

egg shape ~~shape~~




$\hookleftarrow$
moon

If no $\odot$ and ~~response~~ response
equilibrium would be 2 tides / lunar
day. The $\odot$ causes an additional
spring-neap variation.

The tidal potential

Assumptions: all of $\odot$, $\mathbb{A}$, $\oplus$ spherically symmetric, thus $\odot$, $\mathbb{A}$ may be treated as distant pt. masses.

Ellipticity of $\oplus$ would change ht. of theoretical tide by $1/300$ which we neglect.



N.B. α is local
zenith angle
of $\mathbb{A}$

$\underline{O}$: c.o.m. of $\oplus$ (and origin)

$\underline{r}$: pt. of obs. of tide, not necessarily at $\oplus$ surface

r' : dist. from $\mathbb{A}$ or $\odot$ to obs. pt.

$\underline{R}$: posn. of $\mathbb{A}$ or $\odot$

a : radius of $\oplus$

Treat $\mathbb{A}$, $\odot$ separately by principle of superposition.
We consider, e.g., the $\mathbb{A}$.

1. net grav. force of $\mathbb{A}$ on (all of) $\oplus$
is $\underline{f} = G M_{\oplus} M_{\mathbb{A}} \hat{R} / R^2$.

2. acceleration of $\oplus$ in its orbit due to this force

$$\underline{A} = \underline{f} / M_{\oplus} = GM_c \hat{\underline{R}} / R^2$$

N.B. independent of $\underline{r} = \text{obs. pt}$ (all $\leftarrow$ arrows same)

3. grav. pot. of $\mathcal{C}$ at pt. $\underline{r}$

$$\begin{aligned} \phi_c(\underline{r}) &= - \frac{GM_c}{r} \\ &= - \frac{GM_c}{R} \sum_{l=0}^{\infty} \left(\frac{r}{R} \right)^l P_l(\cos \alpha) \\ &\quad \left(\cos \alpha = \hat{\underline{r}} \cdot \hat{\underline{R}} \right) \end{aligned}$$

4. we view tides in an accelerated reference frame, namely on the $\oplus$.
D'Alembert's principle: represent this acceleration by a fictitious force. The resultant is the tidal force. Net tidal force at $\underline{r}$ is:

$$\begin{array}{ccc} -\nabla \phi_c(\underline{r}) & - & \underline{A} \\ \text{variable} \nearrow & & \nwarrow \text{constant} \leftarrow \text{vector} \\ \text{vector} \end{array}$$

$$\begin{aligned} \text{But } -\nabla \phi_c(\underline{r}) - \underline{A} \\ = -\nabla [\phi_c(\underline{r}) + \underline{r} \cdot \underline{A}] \end{aligned}$$

$$\begin{aligned}
 5. \quad \phi_a(\underline{r}) + \underline{r} \cdot \underline{A} &= - \frac{GMa}{R} \left[1 + \frac{r}{R} \cos \alpha \right. \\
 &\quad \left. + \sum_{l=2}^{\infty} \left(\frac{r}{R} \right)^l P_l(\cos \alpha) \right] + \frac{GMa}{R} \frac{r}{R} \cos \alpha \\
 &= - \frac{GMa}{R} - \frac{GMa}{R} \sum_{l=2}^{\infty} \left(\frac{r}{R} \right)^l P_l(\cos \alpha) \\
 &\quad \uparrow \text{const}
 \end{aligned}$$

The const term is the pot. of a at c.o.m. of Φ ; when take ∇ gives rise to no force $\therefore$ is of no interest (the usual arbitrary const. in potential theory).


Note $l=1$ term has been cancelled by A term. This term gives rise to a uniform force throughout Φ , viz. $GM_a \hat{R}/R^2$. But this is precisely the force which keeps Φ accelerating in its Keplerian orbit.

Tidal forces, the resultant, on contrary due to small variations throughout Φ of grav. force/mass exerted by a .

$$c. \text{ Let } V_a(\underline{r}) = - \frac{GMa}{R} \sum_{l=2}^{\infty} \left(\frac{r}{R} \right)^l P_l(\cos \alpha)$$

This called the (inner) tidal potential.

A note on the derivation of the tidal potential:



The total grav. pot. at the point $\bullet$ is:

$$V_{\oplus} \text{ (due to } \oplus) = \frac{GM_a}{r'(t)}$$

Due to its motion relative to inertial space, observer at $\bullet$ also "feels" fictitious forces. The motion at any pt. $\bullet$ can always be regarded as the superposition of motion of c.o.m. (the orbital motion of $\oplus$) + motion w.r.t. c.o.m. (the $\oplus$'s rotation Ω).

The total fictitious force is $-\nabla$ of:

$$-\frac{1}{2}\Omega^2 r^2 \sin^2 \theta + \underline{r \cdot \underline{A}} \quad \begin{array}{l} \text{due to rotation} \\ \text{due to orbital} \end{array}$$


acceleration of c.o.m.

[no change with t on a rigid $\oplus$]

$$\text{But } -\nabla \left[V_{\oplus} - \frac{1}{2}\Omega^2 r^2 \sin^2 \theta \right] = -\nabla U_{\text{geopot.}} = \underline{g}$$

$\underline{f}(\underline{r}) = -\nabla V(\underline{r})$ called tidal force.

Pictorial repr. of $\underline{f}(\underline{r})$ on surface $r=a$ of $\oplus$ is:



Another pt. of view: let $\underline{f}_{\text{tot}}(\underline{r}) \equiv$ ~~force~~ force exerted by moon on a unit mass at $\underline{r}$. Then $\underline{f}(\underline{r}) =$ tidal force $= \underline{f}_{\text{tot}}(\underline{r}) - \underline{f}_{\text{tot}}(\underline{0})$. Tidal force is diff. in force at $\underline{r}$ from force at origin $\equiv$ c.o.m. $\oplus$. (Including the fictitious force due to the orbital acceleration)

Examine relative magnitude of terms in sum (at $\oplus$ surface $r=a$):

$$\underbrace{\left(\frac{a}{R}\right)^2 P_2(\cos \alpha)}_{\text{order 1}} + \underbrace{\left(\frac{a}{R}\right)^3 P_3(\cos \alpha)}_{\text{order 1}} + \dots$$

relative order $\left(\frac{a}{R}\right)$

$$\text{for } \oplus: \frac{a}{R} \sim 1/60$$

$$\text{for } \odot: \frac{a}{R} \sim 1/23,000$$

For $\odot$ can certainly neglect all but $l=2$. For $\oplus$, terms $l > 2$ lead to tides $\sim 1/60$ of $l=2$ tides, can be observed as they occur 3 (and 4, etc.) times per day, but

for moment we neglect them. Then

$$V_a(r) = -\frac{GMa}{R} \left(\frac{r}{R}\right)^2 \left(\frac{3}{2}\cos^2\alpha - \frac{1}{2}\right)$$


lunar tidal potential, similar for $\odot$

What is relative effect of α and $\odot$ in raising tides on $\oplus$?

$$V_a \sim \frac{Ma}{R_a^3}, \quad V_\odot \sim \frac{M_\odot}{R_\odot^3} \quad \text{so}$$

$$a/\odot \sim \frac{M_a/R_a^3}{M_\odot/R_\odot^3} = \frac{\bar{\rho}_a}{\bar{\rho}_\odot} \left(\frac{a_a/R_a}{a_\odot/R_\odot}\right)^3$$

Now a/R is the visible half-aperture in sky.



The visible apertures of α and $\odot$ are very nearly equal (we have eclipses). Thus the effect $\sim$ ratio of densities!

$$\bar{\rho}_a \sim 3.3$$

$$\bar{\rho}_\odot \sim 1.4$$


$$\frac{a}{\odot} \sim \frac{3.3}{1.4} \sim \frac{1}{0.46}$$

a effect a little more than twice $\odot$

Tidal force of a acting up will cause a 90 kg person to lose 10 mg weight (1 part in $\sim 10^7$ $\delta g/g$ change). This $\sim$ weight one drop of sweat.

Question: then how come tides are as large as they are?

Answer: tides are result of tangential forces + cons. of mass (no resistance to tang. motion)



One last question: what is relative effect of $\odot$ and $\oplus$ on lunar tides (on a)?

Solar tides on $a \sim$ solar tides on $\oplus$.

Terr. tides on a :

$$\oplus/\odot \sim \frac{M_{\oplus}/R_{\oplus}^3}{M_{\odot}/R_{\odot}^3} \sim \frac{200}{1}$$

Tides on a dominated by $\oplus$: ~ 200 times


bigger than tides on $\oplus$.

Note: $\mathcal{A}$ always keeps same face toward $\oplus$ (a result of (past) tidal friction, as we shall see). Thus:

monthly ~~tidal period~~ = dominant tidal period on $\mathcal{A}$ due to eccentricity of orbit



The majority of moonquakes (much smaller in energy release or magnitude than large earthquakes) appear to be tidally triggered



No (or only occasional weak) evidence of tidal triggering of quakes on $\oplus$. Two reasons for difference:

1. difference in $\oplus$ & $\mathcal{A}$ interior properties: $\mathcal{A}$ smaller, thicker lithosphere, colder, $\oplus$ convecting, dominated by tectonic strains.
2. tides on $\mathcal{A}$ 200 times bigger than tides on $\oplus$.




Fig. 4. Compressed time-scale records of LP components for two natural seismic events recorded at four seismic stations; X and Y refer to the horizontal component seismometers; Z refers to the vertical component seismometers. Each successive trace deflection of a LP compressed time scale record represents the sum of the absolute values of the differences between 40 successive data samples during consecutive, nonoverlapping 6-s time intervals. Successive deflections are plotted on opposite sides of the zero base line. The relative amplitudes of the signals are indicated by the brackets following each record. Tick marks are 10 min apart, and hours are labeled. The category A₁₈ moonquake zone is located at the eastern border of Mare Serenitatis (see Table 3). For category A moonquakes, the direct shear wave S is prominent on seismograms from the horizontal seismometers. The direct compressional wave P is observed on seismograms of the largest category A moonquakes. The category C meteoroid impacted roughly 500 km north of station 15. The onsets of the direct compressional and shear waves can rarely be identified on LP seismograms of category C events.




Fig. 10. Category A and B seismic activity detected by the Apollo seismic network during the period November 24, 1969, to July 27, 1972. The number of category A and B events detected per day at each station is plotted as a function of time. The time interval covered by each of the stations is given in the figure. The solid triangles indicate times of perigee, and the open triangles indicate times of apogee. The vertical lines below each figure indicate the times of the maximum longitudinal librations. The maximum longitudinal librations occur at times intermediate to apogee and perigee. In general, two peaks in seismic activity are observed at all stations each nodal month at times intermediate to the maximum longitudinal librations. The peaks appear to show no fixed time relationship to apogee and perigee. The dashed curves in each figure represent the number of events detected per nodal month. Points are plotted at times of successive maximum longitudinal librations and each point represents the number of events detected in the preceding and following half-nodal month intervals. A long-period variation (about 206 days) in the number of events detected per month is observed at stations 12, 14, and 15 during 1971 and 1972, but not at station 12 during 1969 and 1970.



The equilibrium tide (Newton)

Consider spherical $\oplus$ with no $\mathcal{L}$ or $\odot$ in sky. Equipotential surfaces are spherical. Consider the surface $r=a$ = radius of $\oplus$.

Let $\phi_0(r)$ be grav. pot. of $\oplus$. On $r=a$, $\phi_0(r) = \phi_0 = -GM_{\oplus}/a = \text{const.}$

Let $g(r) = -r\phi_0'(r)$ grav. ~~acceleration~~ acceleration and let $g \equiv |g(r)|$ on $r=a$ so that $g(r) = -g\hat{r}$ on $r=a$. Then $g = GM_{\oplus}/a^2$.

Now consider $\oplus$ + tide-generating bodies. What will be new, deformed shape of surface with $\text{pot.} = \text{const.} = \phi_0$?



Say pt. $\underline{r}$ moves to $\underline{r} + \underline{h}(r)$. Let $\phi(r) = \text{total pot. including tidal pot. at an arbitrary pt. } \underline{r}$.

$$\phi(\underline{r}) \equiv \phi_0(\underline{r}) + V(\underline{r})$$

On the displaced surface:


$$\begin{aligned} \phi(\underline{r} + \underline{h}) &= \phi_0(\underline{r} + \underline{h}) + V(\underline{r} + \underline{h}) \\ &= \phi_0(\underline{r}) + \underline{h} \cdot \nabla \phi_0(\underline{r}) + \dots \\ &\quad + V(\underline{r}) + \dots \quad \text{where} \end{aligned}$$

... is $O(|\underline{h}|^2)$ which we neglect.

$$= \phi_0(\underline{r}) - \underline{g} \cdot \underline{h} + V(\underline{r})$$

$$= \phi_0(\underline{r}) + g h_r(\underline{r}) + V(\underline{r})$$

now let us denote $h_r \equiv \hat{r} \cdot \underline{h}$
by h



h is the radial displacement (up and down) of equipot. surface.

$$\phi(\underline{r} + \underline{h}) = \phi_0(\underline{r}) + g h(\underline{r}) + V(\underline{r}) + \dots$$

this is $\phi_0 = \text{const.}$ (initial pot. on $r = a$)

we want this to be ϕ_0 too (new total pot. on displaced surface).

Thus on surface of $\oplus$, $r = a$, displacement of equipot. surface is given by

$$h(r) = - \frac{1}{g} V(r) \quad \text{called the equilibrium tide}$$

Note: this same argument we used before to find geoid signal N in terms of anomalous potential V over a mid-ocean ridge.

h , the equilibrium tide: if $\oplus$ were rigid + covered by thin oceans and if tidal potential V changed very slowly so ocean always had time to respond, then $\oplus$ would always be equipotential and the tide would be h . Real tides are dynamical because time scale for changes of $V \approx$ natural resonant periods of ocean basins.

Equilibrium tide still a useful concept: can think of as input to system.

$$h = - \frac{1}{g} V, \quad g = \frac{GM_{\oplus}}{a^2}$$

$$V_a = - \frac{GM_a}{R} \left(\frac{a}{R} \right)^2 P_2(\cos \alpha)$$

Thus
$$h_c = a \frac{M_c}{M_\oplus} \left(\frac{a}{R} \right)^3 P_2(\cos \alpha)$$

Let $\bar{R} \equiv$ mean $\oplus$ distance, a fixed number $\sim 60 \oplus$ radii.

Then

$$h_c(r, t) = K_c \left(\frac{\bar{R}}{R} \right)^3 P_2(\cos \alpha)$$

where $K_c \equiv a \frac{M_c}{M_\oplus} \left(\frac{a}{\bar{R}} \right)^3$

$\xrightarrow{1/81.37}$ $\xrightarrow{(1/60)^3}$

$$= 35.785 \text{ cm}$$

Likewise

$$h_\odot(r, t) = \text{same with } K_\odot$$

$$K_\odot = 16.427 \text{ cm} \sim \frac{1}{2} K_c$$

recall: α is local zenith angle of $\odot$


Note $(\bar{R}/R)^3$ and $P_2(\cos \alpha)$ vary with time.

$\uparrow$ changing distance with time $\uparrow$ changing direction with time.

Both factors are of order 1, so K_c and

$K_{\odot}$ are typical equil. tide heights. Note $K_{\odot} \sim \frac{1}{2} K_a$ as noted already.

Recall that $\alpha(t)$ is angle between $\odot$ or $\ominus$ and local zenith. Consider a standard (geocentric) coord system :



θ, ϕ : colatitude, longitude at station or obs. pt.

$z(t), L(t)$: colatitude, longitude of $\mathcal{C}$ or $\odot$

By spherical trigonometry, or more elegantly by the spherical harmonic addition theorem can be shown that:

$$P_2(\cos \alpha) = \frac{3}{2} \cos^2 \alpha - \frac{1}{2}$$

$$= \left(\frac{3}{2} \cos^2 \theta - \frac{1}{2} \right) \left(\frac{3}{2} \cos^2 \varphi - \frac{1}{2} \right) + \frac{3}{4} \sin 2\theta \sin 2\varphi \cos (L - \phi) + \frac{3}{4} \sin^2 \theta \sin^2 \varphi \cos 2(L - \phi)$$

This the term involving $\alpha(t)$ written in terms of colat. and long. of $\mathcal{A}$ or $\odot$ and station. Can then write equil. tide in form :

$$h_{\mathcal{A}}(\theta, \phi, t) = K_{\mathcal{A}} \left[\frac{3}{2} \cos^2 \theta - \frac{1}{2} \right] \left[\left(\frac{\bar{R}}{R} \right)^3 \left(\frac{3}{2} \cos^2 \mathcal{Z} - \frac{1}{2} \right) \right] \\ + K_{\mathcal{A}} [\sin 2\theta] \left[\frac{3}{4} \left(\frac{\bar{R}}{R} \right)^3 \sin 2\mathcal{Z} \cos(L-\phi) \right] \\ + K_{\mathcal{A}} [\sin^2 \theta] \left[\frac{3}{4} \left(\frac{\bar{R}}{R} \right)^3 \sin^2 \mathcal{Z} \cos 2(L-\phi) \right]$$

equilibrium tide at a point
 θ, ϕ on $\oplus$'s surface as function of time (similarly for $h_{\odot}$)


Sum of 3 terms, each of form

$$K_{\mathcal{A}} [\text{shape of } \oplus\text{'s surface}] [\text{astronomy}(t)]$$


Suppose for moment that $\mathcal{A}$, $\odot$ fixed in space. Then $R(t) = \text{const.}$ and $\mathcal{Z}(t) = \text{const.}$ but $L(t) = -2\pi t / 24 \text{ hrs.}$ due to $\oplus$'s rotation on its axis. This gives the lowest approx. to the tides.

Shows clearly division into 3 principal tidal species.

1. semi-diurnal species : $\cos 2(L-\phi)$
two tidal maxima and minima
per day. Shape or variation
on $\oplus$'s surface $\sim \sin^2 \theta$




2. diurnal species : $\cos(L-\phi)$
one tide / day
shape on $\oplus$ surface $\sim \sin 2\theta$




3. long-term species : $\cos 0(L-\phi) = 1$
no variation at all if $\odot$, $\oplus$
fixed in sky. Actually vary
slowly due to $\odot$, $\oplus$ motion,
largest is fortnightly 2 wk.
period. ~~Shape~~ Shape on
surface $\sim P_2(\cos \theta)$
 $= \frac{3}{2} \cos^2 \theta - \frac{1}{2}$

6 months $\longrightarrow$
for solar
tides



For more accurate picture of tidal variations must consider motions of α , $\odot$ in sky (geocentric view: often adopted, in spite of Copernicus, in discussing positional astronomy, a purely kinematic subject)

Principal features of α , $\odot$ motion:



$\odot$ goes around once per year and α goes around at an inclination of $\sim 5^{\circ}$ to ecliptic once/month.

But due to third body effect of $\odot$ on $\mathcal{A}$ and planetary perturbations (these much smaller but measurable) the orbital parameters change or evolve slowly. Three most rapid secular perturbations together with month and year comprise the:

Five fundamental celestial periods characterizing $\mathcal{A}$ and $\odot$ motion:

1 month = 27.321582 msd, $\mathcal{A}$ declination
1 (tropical) year = 365.242199 msd, $\odot$ declination

8.847 tropical yrs, $\mathcal{A}$ perigee
- 18.613 tropical yrs, regression of lunar node

20,940 yrs, $\odot$ perigee

Next longest fundamental period is precession of equinoxes (Φ 's $\underline{\Omega}$ vector changes position) period $\sim 25,800$ yrs.

How do these celestial periods show up in the description of the equilibrium tide $h(\theta, \phi, t)$?

Consider the three terms we wrote as
[astronomy (t)], viz.

$$[\cdot] = [(\bar{R}/R)^3 (\frac{3}{2} \cos^2 \epsilon - \frac{1}{2})]$$

$$[\cdot] = [\frac{3}{4}(\bar{R}/R)^3 \sin 2\epsilon \cos (L - \phi)]$$

$$[\cdot] = [\frac{3}{4}(\bar{R}/R)^3 \sin^2 \epsilon \cos 2(L - \phi)]$$

A lengthy analytical expansion procedure allows us to write each of these factors in the form:

$$\text{astronomy}(t) = \sum_j C_j \cos(\sigma_j t - m\phi + \theta_j)$$

$\nwarrow$ amplitude $\nearrow$ phase

$m = 0, 1, 2$ long-period,
diurnal, semi-diurnal

$$\sigma_j = m\omega_a + \sum_{i=1}^5 a_i \omega_i, \quad a_i \text{ integers}$$

$\omega_a = \Omega - \omega_1$ so $2\pi/\omega_1 = 1$ mean
lunar day = 24 h 50 m 28 s,
longer than 1 solar day
 $2\pi/\omega_1 = 1$ month, etc., the
five fundamental periods

This a sum of harmonically varying terms. Called harmonic expansion of tidal potential, introduced by Lord Kelvin.


Use of ω_a , lunar day, convenient as lunar tides dominate.

Astronomy gives rise to a modulation of three basic tidal frequencies.

Non-linearity of ~~the~~ celestial mechanics $\rightarrow$ sum and difference frequencies.


Each of the three tidal species split into a number of lines. A line spectrum, but line splitting actually an ∞ process, depends on how much algebra you do and how much resolution.

One speaks of the resolution of any particular development. Results most easily presented in line spectrum form. At 1 cpd resolution, lowest approx., corresponds to a, $\odot$ motionless in space.



At 1 cpm resolution ~~the~~ see fortnightly splitting - each line now called a group

minimum spacing now $2\pi/2w_1 = 13.667$
 days $\sim$ 2 wks (fortnight)



This equivalent
 to assuming
 $C, \odot$ orbits
 are Keplerian
 (unchanging
 elliptical
 elements).

At 1 cpd resolution one sees annual
 splitting of each group — each line now
 called tidal constituent. Minimum
 spacing now $1/2$ years.

Harmonic expansion first suggested by
 Kelvin, carried out by G. Darwin
 $\sim$ 1860's to largest 39 terms, smallest
 $C_j \sim 10^{-3}$ largest. Gave names
 to various constituents, still in use, e.g.

M_2 a semi-diurnal

M_f a fortnightly

S_2 $\odot$ semi-diurnal

A.T. Doodson of Admiralty Office 1921
 Proc. Roy. Soc. carried out expansion
 further to largest 400 C_j , smallest
 $\sim 10^{-4}$ largest. He worked to $\odot$
perigee resolution i.e. found exact speed

of each term, but truncated sum at largest 400.

This a monumental task: 20 years of work on a hand calculator, Doodson's life's work.

His notation (slightly modified) is convenient. Identify a line by its Doodson number:

$$\text{Doodson number} = (m, a_1, a_2, a_3, a_4, a_5)$$

Table of a few of the largest lines:

<u>Doodson</u>	<u>Darwin</u>	<u>period</u>	<u>α or $\odot$</u>	<u>c_j</u>
<u>Long-period:</u>				
020000	Mf	13.661 d	α	.1564
010-100	Mm	27.555 d	α	.0825
<u>Diurnal:</u>				
110000	K_1	1 <u>sidereal</u> day	both	.5305
1-10000	O_1	25.819 hrs.	α	.3769
<u>Semi-diurnal:</u>				
200000	M_2	12.421 hrs.	α	.9081
22-2000	S_2	12 hrs.	$\odot$	.4229

largest α ↑
largest $\odot$ ↑

K_1 at 1 sidereal day = 23h 56m 04s only tidal line due to both α and $\odot$. (also K_2 twice/sid-day)

Show Munk + Cartwright figures of equilibrium spectrum at 1 cpm and 1 cpy resolution. Also plotted are observations of ocean tide at Honolulu, mid-Pacific ocean station. Note noise continuum of meteorological origin.

Also show fortnightly tidal gravity variation measured at South Pole by Louis Slichter.

Final Munk + Cartwright figure shows nodal splitting of S_a and S_{sa} semi-annual and annual lines.

N , the nodal tide, period of 18.6 years.

Note the hint of a pole tide at 14 months, response of the oceans to the Chandler wobble not fixed by $\odot$ or $\oplus$.




FIGURE 2. Honolulu tide spectra at 1 c/y resolution for the low-frequency spherical harmonic P_2^0 . Some 'Darwin' symbols and 'Doodson' constituent numbers are included in the upper panel. Legend is otherwise as in figure 1.




FIGURE 1. Honolulu tide spectra at 1 c/m resolution for spherical harmonics of degree 2. P_2^0 , P_2^1 , P_2^2 refer to tidal species 0, 1, 2 respectively. The upper panel shows the energy of the gravitational equilibrium tide at Honolulu relative to 10^{-4} cm^2 ; some 'Doodson group numbers' are written below the columns. In the next two panels, the observed sea spectrum is designated by the total height of the columns. In the upper of the two, the height of the filled portion designates the energy coherent with the equilibrium tide; in the lower, the height of the unfilled portion designates the noncoherent energy. The left scale refers to energy per column, the right scale to energy per c/d. In panel 4, the filled circles refer to the real part, the open circles to the imaginary part of the admittance. In panel 5, they refer to $|\text{admittance}|$ and phase lead in radians, respectively. The vertical lines show the 95% confidence limits of the circles. The curves represent the admittance functions derived from the convolution process.




FIGURE 3. Honolulu tide spectra at 1 c/y resolution for the diurnal spherical harmonic P_1^1 .




FIGURE 4. Honolulu tide spectra at 1 c/y resolution for the semidiurnal spherical harmonic P_2^2 .




TABLE 2.4.4. Tidal constituents with $C > 0.05$ (from [Bartels, 1957])

Coefficient C	Speed (deg hr ⁻¹) σ	Period 360°/ σ	Symbol (and description)
<i>Long-period tides</i>			
0.2341	0		S_0 (constant solar)
0.5046	0		M_0 (constant lunar)
0.0655	$\omega_4 = 0.00221$	18.613 years	— (nodal to M_0)
0.0729	$2\omega_2 = 0.08214$	182.621 days	Ssa (declinational to S_0)
0.0825	$\omega_1 - \omega_3 = 0.54437$	27.555	Mm (elliptical to M_0)
0.1564	$2\omega_1 = 1.09803$	13.661	Mf (declinational to M_0)
0.0648	$2\omega_1 + \omega_4 = 1.10024$	13.633	— (nodal to Mf)
<i>Diurnal tides</i>			
0.0722	$(\omega_\ell - \omega_1) - (\omega_1 - \omega_3) = 13.39866$	26.868 hours	Q_1 (elliptical to O_1)
0.0710	$(\omega_\ell - \omega_1) - \omega_4 = 13.94083$	25.823	— (nodal to O_1)
0.3769	$(\omega_\ell - \omega_1) = 13.94304$	25.819	O_1 (principal lunar)
0.1755	$(\omega_\odot - \omega_2) = 14.95893$	24.066	P_1 (principal solar)
0.1682	$(\omega_\odot - \omega_2) + 2\omega_2 (= \Omega) = 15.04107$	23.934	K_1^S (declinational to P_1)
0.3623	$(\omega_\ell - \omega_1) + 2\omega_1 (= \Omega) = 15.04107$	23.934	K_1^M (declinational to O_1)
0.0718	$(\omega_\ell + \omega_1) + \omega_4 = 15.04328$	23.931	— (nodal to K_1^M)
<i>Semidiurnal tides</i>			
0.1739	$2\omega_\ell - (\omega_1 - \omega_3) = 28.43973$	12.658 hours	N_2 (elliptical to M_2)
0.9081	$2\omega_\ell = 28.98410$	12.421	M_2 (principal lunar)
0.4229	$2\omega_\odot = 30$	12	S_2 (principal solar)
0.0365	$2\omega_\odot + 2\omega_2 (= 2\Omega) = 30.08214$	11.967	K_2^S (declinational to S_2)
0.0786	$2\omega_\ell + 2\omega_1 (= 2\Omega) = 30.08214$	11.967	K_2^M (declinational to M_2)
<i>Combination tides</i>			
0.5305	$\Omega = 15.04107$	23.934 hours	K_1 (lunisolar declinational)
0.1151	$2\Omega = 30.08214$	11.967	K_2 (lunisolar declinational)




FIGURE 1. Honolulu tide spectra at 1 c/m resolution for spherical harmonics of degree 2. P_2^0 , P_2^1 , P_2^2 refer to tidal species 0, 1, 2 respectively. The upper panel shows the energy of the gravitational equilibrium tide at Honolulu relative to 10^{-4} cm^2 ; some 'Doodson group numbers' are written below the columns. In the next two panels, the observed sea spectrum is designated by the total height of the columns. In the upper of the two, the height of the *filled* portion designates the energy *coherent* with the equilibrium tide; in the lower, the height of the *unfilled* portion designates the *noncoherent* energy. The left scale refers to energy per column, the right scale to energy per c/d. In panel 4, the filled circles refer to the real part, the open circles to the imaginary part of the admittance. In panel 5, they refer to $|\text{admittance}|$ and phase lead in radians, respectively. The vertical lines show the 95% confidence limits of the circles. The curves represent the admittance functions derived from the convolution process.




FIGURE 2. Honolulu tide spectra at 1 c/y resolution for the low-frequency spherical harmonic P_2^0 . Some 'Darwin' symbols and 'Doodson' constituent numbers are included in the upper panel. Legend is otherwise as in figure 1.

(c) *The spectrum at c/y resolution*

Figures 2 to 4 show the same spectra and admittances at greater resolution. Here we see the tidal 'constituents' named by Darwin and other constituents of lesser importance, usually unnamed or allocated to nonlinear 'over-tides'. In figure 2, the constituent labelled MSf is the linear lunar input at $(0 \ 2 \ -2)$, not the $S_2 - M_2$ nonlinear interaction at the same frequency. The coherent energies at 1 and $2c/y$ (Sa and Ssa) were not plotted because they




FIGURE 3. Honolulu tide spectra at $1c/y$ resolution for the diurnal spherical harmonic P_2^1 .

include radiational energy and give anomolous admittances. Separation of gravitational and radiational energy is possible by the method of multiple components. A better procedure is the lumped analysis on which the plotted curves are based. These curves approach a gravitational admittance of $Z = 0.92$ at zero frequency. The reliability of this limit is about the same as that of the estimate at 1 c/m, namely ± 0.24 at the 95 %





FIGURE 4. Honolulu tide spectra at 1 c/y resolution for the semidiurnal spherical harmonic P_2^2 .



GRAVITY TIDE, SOUTH POLE

Tidal prediction


We now have obtained a complete description of cause of tides; the equilibrium tide is a description of tidal forcing or input. We now turn to response of $\oplus$ and oceans to this forcing.

First, a subject of commercial importance, although less so today than in days of sailing ships.

Long before the scientific theorizing, $\exists$ rule of thumb methods of tidal prediction. These considered gainful family possessions, father $\rightarrow$ son, methods undivulged, presumably a form of regression, tides described by a few relevant variables (time, ht. of high and low water, etc.), relate to a few astron. variables (phase, declination of $\odot$, etc.)

Harmonic method developed by Kelvin and Darwin $\sim$ 1860-1870, has a physical basis.

Assumes only that a port's response to tidal input is linear. Let us now denote equil. tide by $\bar{h}(\theta, \phi, t)$



This the reason for harmonic development.
Input at any port

$$\bar{h}(t) = \sum_j \bar{C}_j \cos(\sigma_j t + \bar{\theta}_j)$$

tidal frequencies: $\sigma_j = m\omega_e + \sum_{i=1}^n a_i \omega_i$

↑ amplitude

↑ phase (called $-\phi_{port} + \theta_j$ before).

A linear black box cannot change freq. of input. Resonances can alter amplitude of any harmonic input and resonance + friction can alter phase. One assumes also that $h(t)$ at any port depends only on $\bar{h}(t)$ at the port. Then $h(t)$ must be of form:

$$h(t) = \sum_j C_j \cos(\sigma_j t + \theta_j) \quad *$$

↑ amplitude


↑ phase

↑ tidal frequencies

At any port can determine amplitudes and phases of each line (M_2 , etc.) by analysis of historical tidal records. Then to predict just extrapolate * into future.

Darwin found 39 most important frequencies. By early 1900's method well-established, today in common use. ~~Then~~ Doodson noted that after removal of 39 terms many ~~ports~~ ports still showed strong harmonic residuals. Decided it was worthwhile to find next most important frequencies (went to 400, 20 yrs. of work)

Kelvin invented mechanical analogue machines (predecessor of computer) for harmonic analysis of past tidal records (dial a speed or Doodson number, trace curve, find c_j, θ_j), also invented similar machine for drawing tidal predictions. Samples of both on display in Smithsonian in D.C.



amplitude C_j , phase θ_j , i.e.
 $C_j \cos(\omega_j t + \theta_j)$

Reset C_j, θ_j of lower pulleys for any port
 "... can run off the tides for 1 yr in
 about 4 hrs."

Harmonic method still for most part in use
 today for practical tidal prediction. Uses
 digital electronic computers for analysis.

To what extent do observed tides resemble
 equilibrium? Question is: how close
 are observed C_j, θ_j to $\bar{C}_j, \bar{\theta}_j$?

Admittance defined by $C_j / \bar{C}_j$, resonance
 factor.

Coastal admittances can be up to 10 in
 places.

Even in mid-ocean tides are truly
 a resonant phenomenon.

Example : Honolulu semi-diurnal
admittance varies from $1/2$ to $1 1/2$,
phase lag $\theta_j - \bar{\theta}_j \pm 2$ radians or
more than 90° .

At some ports non-linear interactions
dominate and linear prediction does
not work well.




Figure 7. Tide predicting machine used from 1885 through 1914.




Figure 8. Tide predicting machine used from 1912 through 1965.

Response of the solid Earth : Love numbers

We shall now assume :

1. $\oplus$ has no oceans
2. $\oplus$ is spherically symmetric


Such an $\oplus$ may be described completely for our purposes by three parameters (for any purpose where response is elastic - time scales of a few yrs. or less).

Three parameters :

1. density $\rho(r)$

2. incompressibility or bulk

modulus $\kappa(r)$: a measure of ability to withstand compression.
 Conceptual measurement (defn.):
 Sphere of material, volume V .



pressure P

compression changes volume by ΔV



The bulk modulus is defined by

$$P = \kappa (\Delta V/V)$$

$\uparrow$ pressure applied to outside
 $\uparrow$ fractional change in volume

The more a material can withstand compression the greater is the incompressibility κ

3. rigidity or shear modulus $\mu(r)$:
 a measure of ability to withstand shear. Conceptual measurement



Apply a shearing force F to top of block. Shears by an angle θ . Rigidity is:

$$F/A = 2\mu\theta$$

$\uparrow$ amount of shear
 $\uparrow$ shear force per unit area

For a fluid no rigidity at all
 $\mu = 0$, e.g. air, H_2O ,
 molten Fe of $\oplus$'s core.

Units of κ and μ force/area, dyne/cm².
 Typical values

	κ	μ
H_2O	$0.2 \cdot 10^{11}$ dyne/cm ²	0
granite	$5.5 \cdot 10^{11}$	$2 \cdot 10^{11}$
steel	$17 \cdot 10^{11}$	$8 \cdot 10^{11}$

We shall assume solid $\oplus$ response is
equilibrium since period of $\oplus$ $\ll$ 12 hrs.

Plots of $p(r)$, $\kappa(r)$, $\mu(r)$ for a
 modern $\oplus$ model (PEM) Figures
 6.17 and 6.18 of Stacey, units
 10^{11} Pa or 10^{12} dyne cm⁻².

Features: all increase by about
 one order of magnitude in mantle,
 $\mu = 0$ in fluid ~~mantle~~ core,
 note solid inner core.

Tidal potential deforms $\oplus$. The
surface deformation can be conveniently

precision of observations, the lower mantle and the outer core were each homogeneous and obeyed the Adams-Williamson equation (see also Davies and Dziewonski, 1975). The density profile of one model is shown in Figure 6.17.

Inferences concerning the compositions of the interior zones of the Earth require extrapolation of the model densities to zero pressure.




Figure 6.17. Density profile of Earth model by Dziewonski et al. (1975) (solid line) with corresponding extrapolated zero pressure (and room temperature) density (broken line).

Conventional elasticity theory, which assumes infinitesimal strain, is completely inadequate for this purpose because the elastic constants change dramatically under pressures comparable in magnitude to the elastic moduli. There are first-order changes in density, which can be described only by a finite strain theory. Figure 6.18 is a plot of the elastic moduli within the Earth as functions of pressure. This shows that the variations with pressure are sufficiently regular for simple linear relationships to represent reasonably well the behavior over moderate pressure ranges:

$$K = K_0 + K'_0 P \quad (6.39)$$

$$\mu = \mu_0 + \mu'_0 P \quad (6.40)$$

where subscripts 0 represent zero pressure values. In particular, for the upper part of the lower mantle (from 670 km to about 1500 km according




Figure 6.18. Variations of elastic constants with pressure in the interior of the Earth (model data by Dziewonski et al. 1975—Appendix G).

to the mo

Equation 6
discussion
as Murnag
Equat

which inte



The densit
this relatio
and yields
the adiabatic
ical model
decompress
extrapolati
lieved tha
iron-magne
major disc
lower parts
Dziewonski

Linear
satisfactory
 K_0, K'_0 , whi
relationship
range. Stace
by appealin
ory iron at
coefficient
trapolated
 0.956×10^{11}

to obtain ρ_0
estimate ma
there is no

characterized in terms of 3 parameters, Love numbers, which summarize elastic response of $\oplus$ to tidal ($l=2$) forces.

The surface $r=a$ will be displaced



This the $\oplus$ surface (the ground) being displaced, not equipot. surface.

Let $\hat{r} \cdot \underline{u}(\underline{r}) = u_r(\underline{r}) =$ radial displacement
 Since $\oplus$ a linear system

$$u_r(\underline{r}) = -h \left[\frac{V(\underline{r})}{g} \right]$$

Love number $\nearrow$ $\underbrace{\hspace{1cm}}$ 1 equil. tide

Twice as big a force $\rightarrow$ twice as much displ.

Note $\oplus$ "bulges" where the tidal bulge "bulges".

$h > 0$ for a realizable model.
Actual no. depends on
model $\rho(r)$, $\kappa(r)$, $\mu(r)$
in a complicated way.

For real $\oplus$: $h \sim 0.6$ (0.300 $1066A, B$)

For rigid $\oplus$: $h = 0$, no deformation
at all

For incomp. fluid $\oplus$: $\kappa = \infty$, $\mu = \text{const.}$
 $\rho = \text{const.}$


fluid
 μ, ρ

$$h = 5/2$$

$\kappa = \infty$ not a bad approx for
 $\oplus$ for this application.
even though actually only
about two times μ .

Tangential displacement has two
components. Let θ, ϕ be polar
coords on $r = a$

$\theta = \text{colatitude}$
 $\phi = \text{longitude}$



Then 2 comps. of tang. displ. are:

$$\hat{\theta}: -\ell \left[\frac{1}{g} \frac{\partial V(\theta, \phi)}{\partial \theta} \right]$$

$$\hat{\phi}: -\ell \left[\frac{1}{g} \frac{1}{\sin \theta} \frac{\partial V(\theta, \phi)}{\partial \phi} \right]$$

↑
Love number

Rigid $\oplus$: $\mu = \infty$, $\ell = 0$

Real $\oplus$: $\ell \sim 0.08$ (0.085 1066A, B)

We shall not make use of this Love number further, not introduced by Love but by Shida.

Another quantity of interest is the resulting change in the net grav. pot. at $r=a$ Total pot. is

$$\phi(r) = \phi_0(r) + (1+k)V(r)$$

↑
due to undeformed

$\oplus$ (equal to $-\frac{GM_{\oplus}}{a}$ on $r=a$)

↑ Love number
↑ the applied tidal potential

7

k measures additional pot. due to redistribution of mass of $\oplus$

Rigid $\oplus$: $k = 0$, no redistribution

Real $\oplus$: $k \sim 0.6$ (^{0.3} ~~0.609~~ 1066 A, B)

Incompr. fluid $\oplus$: $k = 3/2$

k and h introduced by A.E.H. Love ~ 1900 .


Together k, l, h characterize equil.

response of spherical $\oplus$ to an applied $l=2$ potential. They are dimensionless numbers, a neat way of summarizing elastic response of $\oplus$.

Their importance : any measurement of tidal response on $\oplus$ surface can be expressed in terms of h, l, k .

Tidal measurements : first example, suppose $\exists$ a grav. potentiometer, could measure grav. potential (actually no such instrument exists but this simplest case to consider).

The instrument is on a surface which moves.



Total potential at $\underline{r} + \underline{u}$ is (Taylor series)

$$\phi(\underline{r} + \underline{u}) = \phi_0(\underline{r}) + \underline{u} \cdot \nabla \phi_0(\underline{r}) + \dots$$

$$+ (1+k) V(\underline{r}) + \dots$$

where ... is $O(u_r^2)$

Now $\underline{u} \cdot \nabla \phi_0 = g u_r(\underline{r})$

$$= - h V(\underline{r})$$


↑ Love number

Thus

$$\phi(\underline{r} + \underline{u}) = \phi_0(\underline{r}) + [1+k-h] V(\underline{r})$$

this pot. on $r=a$
in absence of deformation

Change in potential on displaced surface thus:



$$\delta\phi(\underline{r}, t) = [1+k-h] \\ v(\underline{r}, t)$$



Grav. potentiometer would thus allow measurement of the combination $1+k-h$.

All other measurements on $\oplus$ surface can be expressed in terms of other combs. of Love nos. as we'll see in a moment. First, we note an interesting exploitation of above formula.

Kelvin and Darwin's determination of the rigidity of the $\oplus$:

The height of an equilibrium ocean tide on $\oplus$ acts as a sort of grav. potentiometer.

Tidal forces of $\odot$ and $\ominus$ deform solid $\oplus$ and attract the oceans. If oceans respond in equilibrium they would conform to displaced equipotential



The equilibrium tide on an elastic $\oplus$ is displacement of equipot. surface relative to ocean bottom.

Easy to show that:

direct attraction of $\oplus$, $\ominus$ displaces equipot.

extra pot. due to redistrib. $\oplus$ mass

$$\mathcal{G}(\theta, \phi, t) = - \frac{1}{g} [1 + k - h] V(\theta, \phi, t)$$

$\swarrow$ equilibrium tide
 $\uparrow$ displacement relative to c.o.w. $\oplus$
 $\uparrow$ but ocean bottom moves up

On rigid $\oplus$, $k = h = 0$ and $\mathcal{G} = -\frac{1}{g}V$ as before.

This idea led to first (pre-seismology) measurement of mean rigidity of $\oplus$ as a whole by Kelvin and Darwin in 1880's.


General idea: if $\oplus$ completely rigid then equil. tide $= -\frac{1}{g}V$.
 on a yielding $\oplus$ differs by factor of $1 + k - h$. Kelvin first person to appreciate that tides would deform solid but elastic $\oplus$. Suggested could be employed to find $\mu_{\oplus}$.

Background: accepted view of $\oplus$ at that time: molten interior, thin congealed crust

Reasons:

1. Temp $\uparrow$ as go down in mines.
Simple extrapolation of $(dT/dz)_{surf}$ suggested molten below ~ 100 km. Now known that geothermal ~~gradient~~ gradient decreases with depth.


2. volcanoes



3. important to classical physicists like Kelvin who had read Principia, etc. : $\epsilon \approx \epsilon_{hydro}$ implies $\oplus$ fluid-like.

The diurnal and semi-diurnal tides are not equil. but Kelvin, following Laplace, reasoned that the fortnightly (M_f) tide would be very nearly so since $2 \text{ wks} \gg$ oceanic free periods.

Kelvin used elasticity theory to compute $1+k-h$ for a simple $\oplus$ model (actually he did not employ Love no. notation - introduced later)



radius a

incompressible $\kappa(r) = \infty$

homogeneous $\mu(r) = \mu, \rho(r) = \rho$

uniform rigidity μ , density ρ
 Result : $g = \text{gravity}$

$$k = \frac{3/2}{1 + \frac{19}{2} \frac{\mu}{\rho g a}}, \quad h = \frac{5/2}{1 + \frac{19}{2} \frac{\mu}{\rho g a}}$$

Note for an incomp. fluid $\mu = 0$
 and $k = 3/2$, $h = 5/2$.

Rigid $\oplus$: $\mu = \infty$, $k, h = 0$.

For a fluid $\oplus$: $1 + k - h = 1 + \frac{3}{2} - \frac{5}{2} = 0!$

Makes sense : no ocean bottom, so ht. of tide above ocean bottom zero.

$$1 + k - h = 1 - \frac{1}{1 + \frac{19}{2} \frac{\mu}{\rho g a}}$$

In general :

$$0 \leq 1 + k - h \leq 1$$

$\uparrow$ fluid $\oplus$
 $\uparrow$ rigid $\oplus$

Tide will be reduced on a yielding $\oplus$.
 If 19th cent. view of $\oplus$ correct Mf tide would be very small.

G.H. Darwin undertook analysis of available records : 14 Indian + British ports , 66 years total

Concluded that $1+k+h = 0.7 \pm 0.1$, a reduction of only 30%.

Not an easy measurement : Mf tides are only 1-2 cm in amplitude, g is 0.1564 and shape of $P_2(\cos\theta)$ which is near nodal at mid-latitudes.

$$\frac{1}{1 + \frac{19}{2} \frac{\mu}{\rho g a}} = 0.3 \pm 0.1$$

Knowing $\rho = 5.5 \text{ gm/cm}^3$, $g = 980 \text{ cm/sec}^2$
 $a = 6.371 \cdot 10^8 \text{ cm}$ gives

$$\mu = 8 \cdot 10^{11} \text{ dyne cm}^{-2} \sim \mu_{\text{steel}}$$

A famous result "the $\oplus$ has the rigidity of steel". Very surprising at the time, now only of historical interest.

Seismology has provided us with a much more accurate picture of $\mu(r)$. S waves would not propagate in a fluid.

$\mu(r) = 0$ in fluid core
but overall average for mantle very nearly μ_{steel} , a success for Kelvin to offset his determination of the $\oplus$'s age.


Recent studies of Mf tide at island tidal stations in Pacific show it to be nearly but not exactly "equilibrium" taking defm. of solid $\oplus$ into account.

If admittance defined by:

$$\text{admittance} = \frac{\text{measured amplitude}}{\text{theoretical quasi-static ampl. on yielding } \oplus}$$

then it varies from 0.7 to 1.2 about 20-30% and the phase varies w.r.t. equil. by $\pm 20^\circ$.
Thesis by D. Luther MIT.

These variations due to propagation of



LONGITUDE

Figure 3.4 - Admittance amplitude (a), and phase (b), of the fortnightly (Mf) tide plotted vs. longitude for all stations in the Pacific within 10° degrees of the equator and with more than one year of data. Exceptions are that Buenaventura and La Libertad are not plotted (see text). 95% error bars are indicated.

Figure 3.4a




Figure 3.4b

so-called Rossby waves with speeds much slower than $c = (gH)^{1/2}$

for gravity waves, restoring force for Rossby waves is Coriolis force, a curious beast indeed, of great oceanographical interest.

Reason $\epsilon \approx \epsilon_{\text{hydro}}$ is that behavior of rocks at high temps. and on long time scales is not elastic.




On short time scale (a few years or less) behavior is elastic and θ given by $F/A = 2\mu\theta$ elastic deformation.

On longer time scales even for small loads rock can creep due to motion (glide + climb) of dislocations in x'tals, rate of creep measured by viscosity ν which as we have seen is $\sim 10^{21}$ to 10^{22} for mantle.

Actually, lab experiments on ~~olivine~~ olivine show its creep at high temps. actually governed by a power law relationship, not simple viscosity, $\dot{\epsilon} \propto \tau^3$.

Tidal gravimetry and ocean loading

Best modern measurements of tides of solid $\oplus$ come from tidal gravimeters, measure changes in grav. acceleration on displaced outer surface



what is change in grav. acceleration
 $\equiv$ |grad potential|
 here on $\oplus$ surface?

Derivation slightly more long-winded than for change in potential itself.

Result:

$$\delta g(\underline{r}, t) = \left[1 + h - \frac{3}{2}k \right] \frac{2}{a} V(\underline{r}, t)$$

change in downward
gravitational
acceleration

when $\odot$ or $\ominus$ overhead
 $V(\underline{r}, t)$ is negative

direct
attraction of
 $\odot$ or $\ominus$

lifting of
gravimeter
further from
c.o.m. (note -
sign)

attraction of
tidal
bulge
(note +
sign)

$\odot$ or $\ominus$

$1+h-\frac{3}{2}k$ called the gravimetric factor. It is the combination of Love numbers which can in principle be ~~deduced~~ inferred from tidal gravity measurements.

For a modern $\oplus$ model

$$1+h-\frac{3}{2}k = 1.159 \sim 1.16$$


For many years the goal of $\oplus$ tide research was to measure $1+h-\frac{3}{2}k$ and other Love no. combs. (using other instruments) accurately: use to infer $\rho(r)$, $\kappa(r)$, $\mu(r)$ of $\oplus$.

This program largely unsuccessful. Seismology has given us a much better picture of $\rho(r)$, $\kappa(r)$, $\mu(r)$; can compute Love numbers for realistic $\oplus$ models more accurately than they can be measured.

Gravimeters calibrated to better than 1% but measurements

contaminated by ocean loading effects.

Can plot expected gravity tide in absence of oceans on a phasor plot.



Typically measured $1+h-\frac{3}{2}k$
ranges from 1.10 to 1.25
and phase $\pm 10^\circ - 20^\circ$, i.e.
deviations from theory of order 10%.

Example: superconducting gravimeter
at La Jolla, Calif. and at
Piñon Flat, Calif. (Goodkind
et al.)

Fig. 1 power spectrum of gravity
signal at La Jolla, see
various tidal peaks indicated.




FIG. 1. Power spectrum of gravimeter signal for 100 days at La Jolla.

Note M_m and M_f , etc.

Note also M_3 etc. near 3 cpd
and M_4 etc. near 4 cpd.

These due to neglected terms in
our development of $V(\underline{r}, t)$.

Recall we neglected terms of order

$$\frac{a}{R} = \frac{\text{radius of } \oplus}{\oplus \text{ distance}} \sim \frac{1}{60}$$

We expect M_3 down from M_2 by $1/60$,
 M_4 down $(1/60)^2$.

Amplitude scale is db, 6 db


= factor of 2 so factor of 60

~ 36 db ($2^6 = 64$). This checks.

Fig. 3 shows phasor plots for M_2 and O_1 at La Jolla and Piñon, blown up picture near end of arrow, data $\bullet$, $\blacktriangle$ etc. from different "runs" (need to fill Dewar to maintain liquid He temps several times during a "run").

M_2 lags equilibrium and O_1 leads
it at both stations. Units $1 \mu\text{gal}$
 $= 10^{-6} \text{ gal} \sim 10^{-9} \text{ g}$.

The discrepancy attributed to ocean loading.



Three effects on a gravimeter

1. direct attraction of H_2O below gravimeter
2. depression of ocean floor lowers gravimeter
3. load deforms solid $\oplus$

} these two
largest
both
same
sign.

To correct for these effects need to know ocean tide everywhere, largest effect from nearby oceans, NE Pacific in this case.

↙ we'll discuss these below.

Fig. 2 so-called co-tidal charts

for M_2 and O_1 for this region, one of best known in world, many deep sea capsule drops by group at SIO.

The authors integrated effects of these ocean tides (NEP), added integrated




FIG. 3. Polar plots of ocean loading part of measured and theoretical O_1 and M_2 earth tide at La Jolla and Piñon Flat. The origin (O) is the end point of a vector representing the local theoretical tide of the solid earth with gravimetric factor 1.160 and zero phase lag. The load tide amplitude is represented by the distance from the origin to the observations (⊙, ▲, +, ●) or to the tips of the theoretical load vectors. The phase lag is measured clockwise. At La Jolla ⊙ is run 1, ▲ is run 2, and ● are two 54-day sections of run 2. At Piñon Flat + is run 3 and ● are six 54-day sections of run 3. The theoretical load vectors are calculated for the north-east Pacific Ocean (NEP) and for the 'rest of the world oceans' (WO) which is identical at the two locations. A correction for the height above the local ocean (HC) is made only at La Jolla.





FIG. 2. (a) Modified co-tidal chart for O_1 . (b) Modified co-tidal chart for M_2 .

effects of rest of world's oceans (WO).

Found excellent agreement with observations, observed phase and amplitude to within 0.2 percent.


Pinon $\sim$ 100 miles inland and



loading effects are, as expected, smaller.

At both stations tidal loading effect of order 10%. Mid-continent measurements only affected by $\sim 1\%$. This still too large to allow us to measure some things we would like, e.g. phase lag due to solid $\oplus$ anelasticity, which we expect to be of order $.001^\circ$ to $.01^\circ$ (Zschau) and fluid core resonance at nearly diurnal free wobble period.

Tidal strain measurements affected much more strongly by tidal loading, of order 100% at near coastal stations.



depression of seafloor
causes l to increase.

Also known now that these instruments severely affected by site geometry and local topography. These essentially act to concentrate + modify the global strain field.

There are people now attempting to turn the tide problem around: use grav. tide measurements on land to constrain deep-sea tides in poorly known areas. Each load tide measurement a weighted integral of deep-sea tides, leads to inverse problem.

Co-tidal charts and numerical models:

Recall the ocean tide for a single line ~~line~~ can be written as:

tide as a fun of θ, ϕ and t

$$h(\theta, \phi, t) = C(\theta, \phi) \cos(\sigma t - \Phi(\theta, \phi))$$

amplitude

varies from

place to place

tidal frequency

different for

different lines

phase varies

from

place to place


This conveniently represented by plotting in a so-called co-tidal chart.

Co-tidal lines are lines of constant phase $\Phi(\theta, \phi) = \text{const}$, typically in degrees from 0° to 360° , or in Greenwich hours from 0-12 for M_2 tide. Co-range lines are lines of constant amplitude in cm.

If the M_2 tide were equilibrium the co-tidal chart would look like

$$\bar{h} = 24.25 \sin^2 \theta \cos(\sigma t - 2\phi) \text{ cm.}$$

Taking zero phase at Greenwich.



co-tidal lines are meridians
 ∴ 2 high tides, if one
 on Greenwich then one
 on dateline.

co-range lines are lines of
 latitude.

Actual co-tidal charts for semi-diurnal
 and diurnal much more complicated.
 Originally constructed empirically
 by extrapolation from coastal and
 island observations.


In last few years several
 groups have succeeded in obtaining
 fairly realistic numerical solutions to

Laplace's tidal equations. PDE's which govern response of oceans to tidal input $\bar{h}$, corrected for yielding of solid $\oplus$ below oceans (Laplace assumed $\oplus$ rigid).

Equations depend on knowledge of ocean depth $H(\theta, \phi)$: must be prescribed numerically. Fig. 1 shows $H(\theta, \phi)$ smoothed, note e.g. no Mediterranean. LTE are integro-differential eqns in 3 coupled variables η = tide above ocean floor, u, v = N, E tangential velocity. See eqns (5)-(12) of Accad + Pekeris (1978). Integro-part due to tidal loading and grav. self-attraction.

Solved numerically for M_2, S_2, O_1 , a few others. B.c. no flow into coasts.

Principal feature of co-tidal charts is presence of amphidromal pts. or amphidromes, points where there is zero tidal variation



phase lines
radiate out
like spider legs

Tide progresses around amphidromes either clockwise or counterclockwise. Phase must go from zero to 360° all the way around.

Fig. 9 shows Accad + Pekeris final solution for M_2 tide, note the dozen or so amphidromes, one in ~~the Atlantic~~ N. Atlantic, one in Pacific off coast of La Jolla, etc.

Note "anti-amphidrome" in Indian Ocean, whole basin goes up and down roughly simultaneously.

Fig. 10 shows comparison of phase and amplitude up Pacific Coast.

Fig. 9 shows agreement of theoretical amphidrome in Pacific with observations including several deep-sea capsule drop measurements by IGPP. Tide sweeps up California coast.

Amplitudes disagree 50° - 60° fjord effect local amplification by confined bays.

5

Tide around New Zealand is
interesting.

Tides are high ~ 100 cm a
local maximum in S. Pacific,
but phase goes around ccw
like an amphidrome 180° out
of phase on opposite sides of island.

Figs. 13 + 14 illustrate this in
detail, note coarseness of
discretization.

2. MODIFICATION OF THE TIDAL POTENTIAL DUE TO TIDAL SELF-ATTRACTION AND TO TIDAL LOADING

Let Ω denote the gravitational potential, defined so that the force F per unit mass is given by

$$F = -\text{grad } \Omega. \quad (1)$$

With $\bar{\Omega}$ denoting the astronomical tidal potential, and $\bar{\zeta}$ denoting, in the standard notation (Lamb 1932, p. 332), the equilibrium tide, we have

$$\bar{\Omega} = -g\bar{\zeta}. \quad (2)$$

For the M_2 tide we have

$$\bar{\zeta} = 24.25 \sin^2 \theta e^{i(\sigma t + 2\lambda)} \text{ cm}, \quad (3)$$

where θ and λ denote colatitude and east longitude, and σ is the frequency of M_2 . Let ζ denote the tidal elevation relative to the bottom. If ζ_s is the tidal elevation of the surface relative to the centre of the Earth, and ζ_b the tidal elevation of the bottom relative to the centre of the Earth, then

$$\zeta = \zeta_s - \zeta_b. \quad (4)$$

From this definition of ζ it follows that the equation of continuity retains the Laplacian form

$$\frac{\partial \zeta}{\partial t} = -\frac{1}{a \sin \theta} \left[\frac{\partial}{\partial \theta} (hu \sin \theta) + \frac{\partial}{\partial \lambda} (hv) \right], \quad (5)$$

where u and v denote the components of velocity (assumed uniform throughout the oceanic column), and $h(\theta, \lambda)$ is the variable depth of the ocean. In the presence of a bottom frictional force with components $F_{B\theta}$ and $F_{B\lambda}$, Laplace's tidal equations take the form

$$\frac{\partial u}{\partial t} - 2\omega v \cos \theta = -\frac{g}{a} \frac{\partial \zeta'}{\partial \theta} + \frac{1}{\rho h} F_{B\theta}, \quad (6)$$

$$\frac{\partial v}{\partial t} + 2\omega u \cos \theta = -\frac{g}{a \sin \theta} \frac{\partial \zeta'}{\partial \lambda} + \frac{1}{\rho h} F_{B\lambda}, \quad (7)$$

where ω is the speed of rotation of the Earth. In the original Laplace formulation (Lamb 1932, p. 332), ζ' stood for $(\zeta - \bar{\zeta})$, but this must now be modified to include (1) the effect of the elastic yielding of the Earth to the direct action of the astronomical tidal potential, (2) the effect of the elastic yielding of the Earth to the tidal loading, and (3) the self-attraction of the tides. If $-g\bar{\zeta}$ denotes the change in the gravitational potential due to effects (2) and (3), we have, by conventional theory,

$$\zeta' = \zeta - (1 + k - h) \bar{\zeta} - \bar{\zeta}, \quad (8)$$

where k and h are Love numbers,

$$\xi(\theta, \lambda) = \alpha \sum_n \frac{\epsilon'_n \zeta_n}{(2n+1)} = \frac{\alpha}{4\pi} \int_0^{2\pi} d\lambda' \int_0^\pi \sin \theta' d\theta' \zeta(\theta', \lambda') \sum_n \epsilon'_n P_n(\cos \theta'), \quad (9)$$

$$\epsilon'_n = 1 + k'_n - h'_n, \quad (10)$$

$$\alpha = 3\rho/\bar{\rho} = 0.56, \quad (11)$$

and the axis of the coordinate system (θ', λ') passes through the point (θ, λ) .

Here k'_n and h'_n denote loading Love numbers, $\bar{\rho}$ the mean density of the Earth ($= 5.57 \text{ g cm}^{-3}$), and ζ_n denotes the n th spherical harmonic component of the tide ζ :

$$\zeta = \sum_{n=1}^{\infty} \zeta_n. \quad (12)$$

In (12) we have omitted the term ζ_0 , because it comes out to be only of the order of a millimetre for the two types of boundary conditions which we have used.




FIGURE 1. 1° map. Contour lines of depths of world oceans (in metres).




FIGURE 8. M_2 tide, including effects of self-attraction and of tidal loading. 2° grid, sloping shelf, $h' = 10$ m, $S = 100$ km. —, Cotidal lines (Greenwich hours); ---, orange lines (units: cm).




FIGURE 9. M_2 amphidrome in the northeast Pacific. Observed values (time/amplitude) taken from Irish *et al.* (1971).




FIGURE 10. Comparison of observed with theoretical tides along the northeastern Pacific coast. x, Observed; o, theoretical.




FIGURE 13. The New Zealand coast $P_1 \dots P_6$, and the neighbouring grid-points.




FIGURE 14. Comparison of observed with theoretical tides on the New Zealand coast.
x, Observed; o, theoretical.